



# Mouse IgG Nanoselector Magnetic beads

## Summary

Catalog No	001-101-003
Ligand	Anti-Mouse IgG single domain antibody fragment (VHH, Nanobody)
Bead size	~ 2.8µm
Reactivity	Highly selective for mouse IgG(IgG1, IgG2a, IgG2b, IgG3), no binding to human, rabbit, bovine, horse, sheep and goat IgG.
Binding capacity	High binding capacity, 10 µL slurry bind about 100 µg of recombinant mouse IgG.
Storage	Shipped at ambient temperature. Upon receipt store at 4°C. Stable for 1 year. Do not freeze.
Storage buffer	50 % slurry in PBS containing 20 % Ethanol

## Description

Mouse IgG Nanoselector Magnetic beads have been specifically designed to bind Mouse IgG. Mouse IgG Nanoselector Magnetic beads is based on small high-affinity recombinant alpaca antibody fragments covalently coupled to the surface of Magnetic beads. Mouse IgG Nanoselector Magnetic beads is an ideal tool to isolate or purify Mouse IgG or Mouse IgG Fc-fusion proteins fast and efficiently.

## Background

Mouse IgG Nanoselector Magnetic beads has been specifically designed to bind Mouse IgG. The ligand on the resin is Anti-Mouse IgG single domain antibody fragment(VHH), which is single-domain antibody derived from the variable regions of heavy chain of Camelidae immunoglobulin, also called SdAbs. Compared with protein G, Mouse IgG Nanoselector Magnetic beads has much higher affinity than protein G. Mouse IgG Nanoselector Magnetic beads is effective to pull down Mouse IgG, and 10 µL slurry bind about 100 µg of Mouse IgG.

## Application notes

Protein purification  
Immunoprecipitation (IP)/Co-IP  
Mass spectrometry (MS)

## Benefits

- Highly selective for mouse IgG
- No binding to human, rabbit, bovine, horse, sheep, and goat IgG
- Complex growth media with fetal calf or horse serum can be applied
- Non-animal-derived

## Immunoprecipitation protocol

### Mammalian cell lysis

Note: Harvesting of cells and cell lysis should be performed with ice-cold buffers. We strongly recommend to add protease inhibitors to the Lysis buffer to prevent degradation of your target protein and its binding partners. For one immunoprecipitation reaction, we recommend using ~10<sup>6</sup>- 10<sup>7</sup> cells.

#### 1. Choice of lysis buffer:

\* For cytoplasmic proteins, resuspend the cell pellet in 200 µL ice-cold Lysis buffer by pipetting up and down. Supplement Lysis buffer with protease inhibitor cocktail and 1 mM PMSF (not included).

\* For nuclear/chromatin proteins, resuspend cell pellet in 200 µL ice-cold RIPA buffer supplemented with DNaseI (f.c. 75-150 Kunitz U/mL), MgCl<sub>2</sub> (f.c. 2.5 mM), protease inhibitor cocktail and PMSF(f.c. 1 mM)(not included)

2. Place the tube on ice for 30 min and extensively pipette the suspension every 10 min.
3. Centrifuge cell lysate at 17,000x g for 10 min at +4°C. Transfer cleared lysate (supernatant) to a pre cooled tube and add 300 µL Dilution buffer supplemented with 1 mM PMSF and protease inhibitor cocktail (not included). If required, save 50 µL of diluted lysate for further analysis (input fraction).

### Primary antibody binding(Optional steps)

1. take supernatant of the cell lysis into another tube and then add primary antibody (2ug-4ug/sample).
2. rotate at +4°C overnight.

This product is for research use only and is not approved for use in humans or in clinical



#### Beads equilibration

1. Resuspend the beads by gently pipetting up and down or by inverting the tube. Do not vortex the beads!
2. Transfer 25  $\mu$ L of bead slurry into a 1.5 mL reaction tube.
3. Add 500  $\mu$ L ice-cold Dilution buffer.
4. Separate the beads with a magnet until the supernatant is clear.
5. Discard the supernatant.

#### Protein binding

1. Add diluted lysate to the equilibrated beads.
2. Rotate end-over-end for 1 hour at +4°C.

#### Washing

1. Separate the beads with a magnet until the supernatant is clear.
  2. If required, save 50  $\mu$ L of supernatant for further analysis(-flow-through/non-bound fraction).
  3. Discard remaining supernatant.
  4. Resuspend beads in 500  $\mu$ L Wash buffer.
  5. Separate the beads with a magnet until the supernatant is clear. Discard the remaining supernatant.
  6. Repeat this step at least twice.
  7. During the last washing step, transfer the beads to a new tube.
- Optional: To increase stringency of the Wash buffer, test various salt concentrations e.g. 150 mM - 500 mM, and/or add a non-ionic detergent e.g. Triton™ X-100.

#### Elution with 2x SDS-sample buffer

1. Remove the remaining supernatant.
2. Resuspend beads in 80  $\mu$ L 2x SDS-sample buffer.
3. Boil beads for 5 min at +95°C to dissociate immunocomplexes from beads.
4. Separate the beads with a magnet.
5. Analyze the supernatant in SDS-PAGE.

#### Elution with Glycine-elution buffer

1. Remove the remaining supernatant.
2. Add 50–100  $\mu$ L Glycine-elution buffer and constantly pipette up and down for 30 - 60 sec at +4°C.
3. Separate the beads with a magnet until the supernatant is clear..
4. Transfer the supernatant to a new tube.
5. Immediately neutralize the eluate fraction with Neutralization buffer.
6. Repeat this step at least once to increase elution efficiency .

### Suggested buffer compositions

Buffer	Composition
Lysis buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.5 % NP40
RIPA buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.1 % SDS, 1 % Triton™ X-100, 1 % deoxycholate
Dilution/Wash buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA
2x SDS-sample buffer	120 mM Tris/Cl pH 6.8, 20 % glycerol, 4 % SDS, 0.04 % bromophenol blue, 10 % $\beta$ -mercaptoethanol
Glycine-elution buffer	200 mM glycine pH 2.0
Neutralization buffer	1 M Tris pH 10.4