

Anti-Mouse IgG kappa, AlpSdAbs[®] VHH(Cy3B ×4)

Summary

Code	001-106-015
Immunogen	Fab region of Mouse IgG
Host	Alpaca pacous
Isotype	VHH domain of alpaca IgG2b/2c
Conjugate	Cy3B(Ex=560nm, Em=571nm), 2 moles Cy3B per mole VHH
Specificity	Mouse IgG kappa chain
Cross-Reactivity	No cross-reactivity with rabbit, human, cynomolgus, rat, goat IgG
Purity	Recombinant Expression and Affinity purified
Concentration	1mg/ml
Formation	Liquid, 10mM PBS (pH 7.5), 0.05% sucrose, 0.1% trehalose, 0.01% proclin300, 50% Glycerol
Storage	Store at -20 °C(Avoid freeze / thaw cycles) , protect from light

Description

Anti-Mouse IgG kappa, AlpSdAbs[®] VHH(Cy3B ×4) is designed for detecting mouse IgG in super-resolution microscopy techniques specifically. Anti-Mouse IgG kappa, AlpSdAbs[®] VHH(Cy3B ×4) is based on monoclonal, recombinant single domain antibody to mouse IgG kappa chain coupled to Cy3B. Based on immunoelectrophoresis and/or ELISA, Anti-Mouse IgG kappa, AlpSdAbs[®] VHH(Cy3B ×4) reacts with the mouse IgG kappa chain selectively, no

Background

VHH are single-domain antibodies derived from the variable regions of heavy chain of Camelidae immunoglobulin. The size of VHH is extremely small(<15KDa) compared to other forms of antibody fragment, which significantly increase the permeability of VHH. Thus VHH is considered of great value for research, diagnostics and therapeutics.

Standard immunodetection approaches use typically a primary antibody (1.Ab) which binds the protein of interest (POI) and a secondary antibody (2.Ab) that binds to the 1.Ab and carries a detection element. The complex formed by the primary antibody and the secondary antibody (1.Ab-2.Ab) is widely used because it is a cost effective and flexible approach since only the 2.Abs need to be coupled to the detection element. However, the use of this complex carries some relevant limitations. The 1.Ab-2.Ab can measure up to 30 nm, leading to a large distance between the targeted molecule and the detection element, causing the so called "linkage" or "displacement" error. While this might not influence the results in some applications (e.g. epifluorescence, ELISA or FACS), it is of major relevance for super-resolution microscopy techniques where the localization precision can be as high as 1 nm. The linkage error can be reduced by using directly labelled small affinity probes like camelid single domain antibodies (sdAbs) also known as nanobodies (Nbs), which have sizes below 3 nm.

Benefits

High lot-to-lot consistency
 Increased sensitivity and higher affinity
 Animal-free production

Application notes

IF/Super-resolution microscopy (iStorm, etc)
 The recommended dilution ratio is 1:100-1:1000.

Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically.

This product is for research use only and is not approved for use in humans or in clinical