

# c-Myc Peptide

## Properties

Product Name	c-Myc Peptide
Code	APK052
Peptide	EQKLISEEDL
Assay	≥95% (HPLC/MS)
Form	Lyophilized powder
Storage Temp	2-8°C
Preparation Note	Dissolve in TBS as a stock solution(Storage at -20°C) at a concentration of 5 mg/ml.

**Note:** TBS(50 mMTris-HCl, pH 7.4, with 150 mM NaCl)

## Description

The c-Myc Peptide is a synthetic peptide of 10 amino acid residue(EQKLISEEDL).

## Benefits

After competitive elution of the c-Myc tag fusion expression protein from this product, the eluted sample does not contain the light and heavy chains of the traditional Anti-myc tag antibody, effectively solving the interference issue of light and heavy chains during Western detection after immunoprecipitation. In conventional immunoprecipitation, the c-Myc tag fusion expression protein is usually released into the solution by boiling with the addition of loading buffer, but at this time, the light and heavy chains of the Anti-myc tag antibody also enter the solution and are likely to interfere with subsequent Western detection. The most important thing about peptide elution is to maintain the activity of the elution product.

## Application

Recommended working concentration is 100-400µg/ml for elute 3×Flag fusion proteins from the c-Myc affinity gel(Myc tag Nanoselector Agarose, Myc tag Nanoselector Magnetic beads or other Anti-Myc affinity gel).

Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically.

This product is for research use only and is not approved for use in humans or in clinical